CTS-6 Evaluation Tool

>12 = 0.80 probably of carpal tunnel syndrome >5 = 0.25 probably of carpal tunnel syndrome

EastBayH

The Value Added by Electrodiagnostic Testing in the Diagnosis of Carpal Tunnel Syndrome Brent Graham J Bone Joint Surg Am. 2008;90:2587-2593

Symptoms and History	
1. Numbness predominantly or exclusively in the median nerve territory	(3.5)
Sensory symptoms are mostly in the thumb, index, middle and/or ring fingers	
2. Nocturnal numbness	(4)
Symptoms are predominantly the patient sleep; numbness wakes patient from sleep	
Physical examination	
3. Thenar atrophy and/or weakness	(5)
The bulk the thenar area is reduced or where manual motor testing shows strength of grade 4 less	
4. Positive Phalen's test	(5)
Flexion of the wrist reproduces her worsened symptoms of numbness in the median nerve territory	
5. Loss of 2 point discrimination	(4.5)
Failure to discriminate 2 points held 5 mm or less apart from one another, in the median innervated dig	its
6. Positive Tinel sign	(4)
Light tapping over the median nerve at the level of the carpal tunnel causing radiating paraesthesias	(-/
Total	(26)
	(=0)

*Disclaimer: This tool is modified from published literature validating the use of the CTS-6. It is not intended to be used for the diagnosis of any medical condition without evaluation by your physician. The diagnosis and treatment of any medical condition should be performed by a trained physician.

